

MERIT beam request


A.Fabich
CERN AB-ATB

<http://cern.ch/proj-hiptarget>

Collaboration meeting, October 2005




Contents



- Summary of previous actions
 - Pulse list
 - Beam request to the APC
- Pulse list
- Response of APC
- Next steps



Available Information



APC: Accelerator Performance Committee


- Platform to discuss all issues relevant to CERN accelerators
- <http://ab-div.web.cern.ch/ab-div/Meetings/APC/Welcome.html>

Following the approval of MERIT:

- Presentation of beam request to APC (June 2005)
 - http://proj-hiptarget.web.cern.ch/proj-hiptarget/default/Documents/subsystems/ProtonBeam/APC_Jun05_AFabich.ppt
 - APC asked for a pulse list as follow-up action:
- MERIT pulse list: <http://proj-hiptarget.web.cern.ch/proj-hiptarget/redir.asp?short=pulselist> (July 2005)
- APC responded on feasibility of configurations



MERIT pulse list




Version in June was a suggestion for discussion:

- 80 different configurations
 - Each two pulses
 - Total: 160 pulses = 4.4×10^{15} protons on TARGET
 - Remember: limited to 3×10^{15} p.o.t. by RP safety
 - varying intensity, time structure, displacement, spot size and solenoid field
 - Solenoid field not in the responsibility of PS
 - About 25 different proton beam configurations



MERIT pulse list (ff)



#	B-field	momentum	protons/bunch	bpp	buckets	dx	dy	spot rms	scan
	[T]	[GeV/c]	[*10^12 p+]			[mm]	[mm]	[mm]	
1	0	24	1	4	1-2-3-4	0	0	1.2	
2	0	24	2	4	1-2-3-4	0	0	1.2	
3	0	24	3	4	1-2-3-4	0	0	1.2	
4	0	24	4	4	1-2-3-4	0	0	1.2	
5	0	24	5	4	1-2-3-4	0	0	1.2	
6	0	24	6	4	1-2-3-4	0	0	1.2	
7	0	24	7	4	1-2-3-4	0	0	1.2	
8	3	24	7	4	1-2-3-4	0	0	1.2	
9	6	24	7	4	1-2-3-4	0	0	1.2	
10	9	24	7	4	1-2-3-4	0	0	1.2	
11	12	24	7	4	1-2-3-4	0	0	1.2	
12	15	24	7	4	1-2-3-4	0	0	1.2	
13	-5	24	7	4	1-2-3-4	0	0	1.2	
14	-10	24	7	4	1-2-3-4	0	0	1.2	
15	-15	24	7	4	1-2-3-4	0	0	1.2	
16	3	24	4	4	1-2-3-4	0	0	1.2	
17	6	24	4	4	1-2-3-4	0	0	1.2	
18	9	24	4	4	1-2-3-4	0	0	1.2	
19	12	24	4	4	1-2-3-4	0	0	1.2	
20	15	24	4	4	1-2-3-4	0	0	1.2	
21	15	24	7	4	1-2-3-4	-7	0	1.2	
22	15	24	7	4	1-2-3-4	-5	0	1.2	
23	15	24	7	4	1-2-3-4	-3	0	1.2	
24	15	24	7	4	1-2-3-4	3	0	1.2	
25	15	24	7	4	1-2-3-4	5	0	1.2	
26	15	24	7	4	1-2-3-4	7	0	1.2	
27	15	24	7	4	1-2-3-4	0	-7	1.2	
28	15	24	7	4	1-2-3-4	0	-5	1.2	
29	15	24	7	4	1-2-3-4	0	-3	1.2	
30	15	24	7	4	1-2-3-4	0	3	1.2	
31	15	24	7	4	1-2-3-4	0	5	1.2	
32	15	24	7	4	1-2-3-4	0	7	1.2	
33	0	24	7	4	1-2-3-6	0	0	1.2	
34	15	24	7	4	1-2-3-6	0	0	1.2	
35	0	24	7	4	1-2-3-8	0	0	1.2	
36	15	24	7	4	1-2-3-8	0	0	1.2	




#	B-field	momentum	protons/bunch	bpp	buckets	dx	dy	spot rms	size
	[T]	[GeV/c]	[*10^12 p+]			[mm]	[mm]	[mm]	
37	0	14	7	4	1-2-3-4	0	0	minimum	
38	15	14	7	4	1-2-3-4	0	0	minimum	
39	0	14	7	4	1-2-3-7	0	0	minimum	
40	15	14	7	4	1-2-3-7	0	0	minimum	
41	0	14	7	4	1-2-3-10	0	0	minimum	
42	15	14	7	4	1-2-3-10	0	0	minimum	
43	0	14	7	4	1-2-3-12	0	0	minimum	
44	15	14	7	4	1-2-3-12	0	0	minimum	
45	0	14	7	4	1-2-3-14	0	0	minimum	
46	15	14	7	4	1-2-3-14	0	0	minimum	
47	0	14	7	4	1-2-3-16	0	0	minimum	
48	15	14	7	4	1-2-3-16	0	0	minimum	
49	0	14	7	4	1-2-3-20	0	0	minimum	
50	15	14	7	4	1-2-3-20	0	0	minimum	
51	0	14	7	4	1-2-3-28	0	0	minimum	
52	15	14	7	4	1-2-3-28	0	0	minimum	
53	0	14	7	4	1-2-3-40	0	0	minimum	
54	15	14	7	4	1-2-3-40	0	0	minimum	
55	0	14	7	4	1-2-3-400	0	0	minimum	
56	15	14	7	4	1-2-3-400	0	0	minimum	
57	0	24	7	8	2 bunches every 20 ms	0	0	1.2	
58	8	24	7	8	2 bunches every 20 ms	0	0	1.2	
59	15	24	7	4	2 bunches every 20 ms	0	0	1.2	
60	0	24	7	8	4 bunches every 20 ms	0	0	1.2	
61	8	24	7	8	4 bunches every 20 ms	0	0	1.2	
62	15	24	7	4	4 bunches every 20 ms	0	0	1.2	
63	0	24	7	4	1-2-5-6	0	0	1.2	
64	15	24	7	4	1-2-5-6	0	0	1.2	
65	0	24	7	4	1-2-7-8	0	0	1.2	
66	15	24	7	4	1-2-7-8	0	0	1.2	
67	0	14	7	4	1-2-9-10	0	0	minimum	
68	15	14	7	4	1-2-9-10	0	0	minimum	
69	0	14	7	4	1-2-11-12	0	0	minimum	
70	15	14	7	4	1-2-11-12	0	0	minimum	
71	0	14	7	4	1-2-13-14	0	0	minimum	
72	15	14	7	4	1-2-13-14	0	0	minimum	
73	0	14	7	4	1-2-15-16	0	0	minimum	
74	15	14	7	4	1-2-15-16	0	0	minimum	
75	0	24	7	4	1-2-3-4	0	0	1.4	
76	0	24	7	4	1-2-3-4	0	0	1.6	
77	0	24	7	4	1-2-3-4	0	0	1.8	
78	0	24	7	4	1-2-3-4	0	0	2	
79	0	24	7	4	1-2-3-4	0	0	2.3	
80	0	24	7	4	1-2-3-4	0	0	2.6	

pump-probe method

rep rate


pump-probe method 2

spot size





Summary of pulse list



- Counting only PS beam parameter changes
 - About 30 different beam configurations: varying
 - Intensity
 - Beam displacement on target (hor./ vert.)
 - Pump-probe method
 - 50-Hz operation
 - Spot size
- APC statement: two “surprises”
 - Intensity limited to 4×10^{12} /bunch (harmonic 8)
 - 50 Hz operation not possible
 - Kicker can do it, but not the septum

Comments to the configuration list


- # The **majority** of the bunch configurations can be made using correct **PS Booster synchronisation** and by timing the **PS extraction kicker** precisely.
- # The **double batch** extraction separated by 20 ms **cannot be done** (6 cases).
- # This means that **10 out of 80** cases cannot be done.
- # The **intensity** that can be guaranteed is **4E12** protons per bunch.
- # The beam spot is based on optics calculations made by the collaboration. The collaboration is responsible for the result. We will provide the initial beam parameters.
- # Some MD time, in order to increase the intensity per bunch, can be allocated end 2006.

Proposals and Conclusion:

- For obtaining higher intensities:
 - One should look into using **higher harmonics** (h_{16} instead of h_8), using more bunches in same time slot.
 - One should consider the possibility of **double batch injection** in combination with the above.
 - One should try **not** to do the experiment **within the two months after the machine start-up**.
- We will **make another iteration** on the configuration list with A. Fabich, taking some of the previously mentioned proposals into account in order to decide the MD subject(s).



How to overcome restrictions?



■ Intensity?


- Consider running PS in harmonic 16
 - increases intensity to 5×10^{12} p⁺ per 1/8 of PS
 - Beam structure changes to double number of bunches in same time interval

■ 50 Hz operation

- Operate at 14 GeV/c only (pulse #57-62)
- Like pump-probe method, but extend bunch-to-bunch distance extensively (milliseconds)



Next steps



MERIT pulse list:

- Next iteration of proposal process
- Incorporate suggestions of APC
- Finalize list of requested beam configurations

- Estimate MD time needed

- PS-SPS coordinator was present at APC
 - Re-discuss with him possible dates of beam time